

THE NANOBIOLOGICAL MENACE OF BLACK HELICOPTERS

RESEARCH BY LYLE ZAPATO, ZPi LABS, REPUBLIC OF CASCADIA

WHAT ARE BLACK HELICOPTERS?

Black Helicopters (BHs) are autonomous agents—lifeforms—created by the **New World Order (NWO)** using top-secret **nanobiotechnology**.

BH REPRODUCTION, DEVELOPMENT, & LIFESTYLE


The BH reproductive cycle starts with **seed crystals (SC)** that are injected into biological hosts—typically cattle or humans—by either human technicians or other BHs. SCs may also be sprayed from aircraft (so-called "**chemtrails**"). The inoculated SC uses **nanobiotechnology** and compounds gathered from the host's organs to grow as it enters the blood stream and travels to the intestines. Once there, it develops into a **Microscopic Black Helicopter (MBH)**.

MBHs will either exit the body through the skin or stay in the host and grow, reaching about 20-30 cm long in a few weeks before bursting forth from the host's guts. This **artificial live birth** kills the host and is the source of **cattle mutilations**. Bursting usually occurs at night. The new BH's props unfold and harden before it flies off to gather minerals and proteins from its environment so it can grow to full mature size (up to 18 meters long.)

Macroscopic BHs not yet their mature size are called **Juvenile Black Helicopters (JBH)**. If discovered, JBHs may eliminate any witnesses or use their psychotron organs grown from environmental aluminum compounds to cause short- to mid-term memory loss (known as "**lost time**") in observers before escaping. Smaller JBHs may camouflage themselves as pigeons, squirrels, and other woodland or urban creatures by covering themselves with the outer casings of animals they killed for organic building materials.

A BH can inject SCs when it has reached the MBH stage of development. When its injector grows too big to puncture flesh without causing hemorrhaging, it is instead used as a projectile weapon. A MBH that has exited its host before bursting (0.25 to 1 cm long) is called an **Extracorporeal Microscopic Black Helicopter (EMBH)**. Often the source of seed injections, they are also called **reproductives** or **seed vectors**. They are almost undetectable and can act as spies. **Reinfections** occur when an EMBH re-enters a host through the ear, burrows to the skull base, and implants itself in the central nervous system, allowing it to affect or even control the host's actions. This is dangerous, since the MBH—now called a **Parasitic Black Helicopter (PBH)**—can do permanent brain damage and may self-destruct to avoid detection, causing what is often erroneously diagnosed as **spontaneous human combustion**.

BLACK HELICOPTER LIFECYCLE



NOTE: Infected cattle can also be under control of their PBH, causing them to become NWO agents (misleadingly termed **Mad Cows** in the NWO-controlled press after the need to cover up any evidence of their conspiracy).

CONCLUSIONS

Black Helicopters represent a **SIGNIFICANT NANOBIOLOGICAL MENACE** to the future of Humanity far beyond **grey goo** or **uppity sentient nanobots** since BHs were **EXPLICITLY DESIGNED FOR EVIL**. Nanotech researchers should stop wasting their time on nanotoothpaste and very small guitars, and instead focus on building counter-measures such as **nanosurface-to-air missiles** and the **nanofreedom fighters** to wield them.